GLOBAL MARK			
CLEAR PASS	BORDERLINE PASS	BORDERLINE FAIL	FAIL

## Criteria

Criteria	Т
Introduces Self, confirms patient identify, Washes hands,	2, 1
PPE	1
Confirms team competencies	1
Explains Plan	
Summons appropriate help	
A – applies 15L O2	
B – quick assessment	
C – Asks for 12 lead, notes BP, and plethora, cannulates, sends bloods	
inc Troponin	
D – GCS 15, Checks BM	
E – ECG shows acute MI, knows territory	
V1-2 Septal - LAD	
V3-4 Anterior - LAD	
V5-6 Apical – LAD or RCA	
I, aVL Lateral LCx	
II aVF, III Inferior RCA	
Treats things appropriately	
Asks for Morphine, Nitrates, Aspirin, Prasugel	
Verbalises need for primary PCI	
Recognises cardiac arrest	
Starts CPR 30:2	
Manages airway – BVM and adjuncts	
Applies defibrillator pads if not already done so	
Recognises rhythm	
Delivers shock safely	
Considers Reversible Causes	
Hypoxia, Hypovolaemia, Hypothermia, Hyperkalaemia	
Thrombosis, Tamponade, Toxins, Tensions Pneumothorax	
Continues CPR	
Considers ECHO in life support	
Minimizes interruptions to CPR	
Utilizes ETCO2	
Discusses thrombolysis if there is going to be a long wait for PCI	
Alteplase 15mg IV bolus + 50mg over 30 mins + further infusion	
(max dose 100mg 90 mins).	
Deals with less than helpful cardiology SpR	
Plans for safe transfer of patient to cath lab/ICU	
Thanks team	
	Tom Birchar 2010