

PEPSE

GLOBAL MARK			
CLEAR PASS	BORDERLINE PASS	BORDERLINE FAIL	FAIL

Criteria

Introduces Self, confirms patient identify, Washes hands,	2, 1
Offers analgesia, ensures comfort of patient	2
Confirms nature of the problem	1
Is non-judgemental	
Ascertains exactly what kind of exposure Anal Vaginal Oral Multiple partners Any bleeding?	
Ascertains exactly what kind of partner Male/Female Ethnicity BBV status (if known) IVDU	
Ascertains when the exposure was	
Makes an assessment of risk based on known information	
Makes reference to using BASH guidelines to calculate the risk formerly, but essentially – ANAL, VAGINAL, SHARING OF NEEDLES → PEP if HIV poorly controlled or high risk group If HIV viral load <200 or Not high risk group – NO PEP	

Table 3. Summary table of PEPSE prescribing recommendations.

	Source HIV status			
	HIV-positive		Unknown HIV status	
	HIV VL unknown/ detectable (>200 copies/mL)	HIV VL undetectable (<200 copies/mL)	From high prevalence country/risk-group (e.g. MSM) ^a	From low prevalence country/group
Receptive anal sex	Recommend	Not recommended ^b Provided source has confirmed HIV VL < 200 c/mL for > six months	Recommend	Not recommended
Insertive anal sex	Recommend	Not recommended	Consider ^c	Not recommended
Receptive vaginal sex	Recommend	Not recommended	Consider ^c	Not recommended
Insertive vaginal sex	Consider ^d	Not recommended	Consider ^c	Not recommended
Fellatio with ejaculation ^e	Not recommended	Not recommended	Not recommended	Not recommended
Fellatio without ejaculation ^e	Not recommended	Not recommended	Not recommended	Not recommended
Splash of semen into eye	Not recommended	Not recommended	Not recommended	Not recommended
Cunnilingus	Not recommended	Not recommended	Not recommended	Not recommended
Sharing of injecting equipment ^f	Recommended	Not recommended	Consider	Not recommended
Human bite ^g	Not recommended	Not recommended	Not recommended	Not recommended
Needlestick from a dis- carded needle in the community			Not recommended	Not recommended