

Sexually Transmitted infection

GLOBAL MARK			
CLEAR PASS	BORDERLINE PASS	BORDERLINE FAIL	FAIL

Criteria

Introduces Self, confirms patient identify, Washes hands,	2, 1
Offers analgesia, ensures comfort of patient	2
Uses open questions to find out what happened Explains questions are broad and can be a little difficult to answer sometimes. Stresses confidentiality	2
Explores current symptoms <ul style="list-style-type: none"> • Urethral discharge • Genital ulceration • Vaginal discharge/bleeding • Deep dyspareunia/abdo/pelvic pain 	
Explores contraception practice <ul style="list-style-type: none"> • Type • Compliance 	
Gets specifics <ul style="list-style-type: none"> • Last sexual intercourse • Risk assessment of partner for BBV and STIs • Type of last sexual intercourse (vaginal, anal, oral) • Number of partners 	
Past sexual history <ul style="list-style-type: none"> • Ever been paid for sex • Ever had sex with a man • Explores for intimate partner violence and sexual assault • Considers child and adult protection issues 	
Past Gyne history <ul style="list-style-type: none"> • Pregnancy and miscarriage • Cervical smear status, and HPV vaccination status 	
Alcohol and Recreational Drug History	
Smoking status	
Drugs and Allergies	
Social History Long term partner Children	
PMH	
Considers <ul style="list-style-type: none"> • PEP (Truvada and Raltegravir) • Accelerated Hep B vaccination 	
Explains <ul style="list-style-type: none"> • Onwards referral to STI clinic as appropriate 	

