Sexual Assault

GLOBAL MARK				
CLEAR PASS	BORDERLINE PASS	BORDERLINE FAIL	FAIL	

Criteria

Criteria	
Introduces Self, confirms patient identify, Washes hands,	2, 1
Offers analgesia, ensures comfort of patient	2
Uses open questions to find out what happened	2
Finds out what kind of assault	2
Oral rape	
Vaginal rape	
Anal rape	
Digital or penile penetration	
Asks about injuries	2
Bleeding	
Genital injuries	
Other injuries	
Offers support	2
Explains the patient is the victim of a crime	
Offers police involvement in a non-coercive way	
Offers SARC involvement	
Explains role of SARC	2
Forensic examination	
Samples kept for 7 years	
Offers morning after pill	2
Explores need for PEP	2
Identity of attacker (known HIV, BBV, African)	
Talks about using BASH guidelines for PEP	
If unknown and no risk factors – risk is small	
If there is bleeding risk is higher	
If multiple rapists, anal, bites, wounds risk is higher	
If HIV+VE or IVDU – risk is still small but higher	
Explores wider safety issues	2
Child Protection	
Vulnerable adults	
Knives/Guns involved	
Explains will need to examine for injuries	2
But examination will not be a forensic examination	
Double gloved for examination	
Tries not to cross contaminate different body areas	
Offers Prophylactic antibiotics for STI	2
Cefixime 400mg Azithromycine 1g, Metronidazole 2g	
Explains SARC referral	2
Asks patient to not brush teeth	
If clothes are removed place in a sealed bag	

Suggests attending GUM clinic within 2 weeks of assault for further investigations	
Explores safe discharge planning	2
Gives advice about rape crisis line, victim support, havens	
Total	

Tom Bircher 2019